

GnRH analogues may increase endometrial Hoxa10 promoter methylation and affect endometrial receptivity

FEI LI, MING ZHANG, YUANZHEN ZHANG, TENG LIU and XINLAN QU

Department of Obstetrics and Gynecology, Reproductive Medicine Center, Zhongnan Hospital of Wuhan University, Wuhan, Hubei 430071, P.R. China

Received October 13, 2013; Accepted May 2, 2014

DOI: 10.3892/mmr.2014.2680

Abstract. The present study aimed to investigate whether gonadotropin-releasing hormone analogues (GnRH-as), including GnRH agonists and antagonists, affect endometrial homeobox (Hox) a10 DNA methylation during the implantation window in mice. GnRH analogue mouse models were used and were treated with either human menopausal gonadotropin (HMG) and a GnRH agonist or HMG and a GnRH antagonist. Uterus samples were collected 48 h after GnRH analogue treatment or ovulation. Bisulfite sequencing polymerase chain reaction (PCR), quantitative-PCR and western blot analysis were performed to assess Hoxa10 and integrin β 3 expression. Scanning electron microscope analyses were conducted to analyze pinopode development. Compared with the natural cycle control mice, mice in the GnRH analogue groups were found to exhibit increased levels of methylation at the Hoxa10 promoter, decreased Hoxa10 mRNA and protein expression and disrupted pinopode development. These findings suggest that GnRH-as may be associated with altered Hoxa10 DNA methylation, thus GnRH-as may affect uterine Hoxa10 expression and endometrial receptivity.

Introduction

Controlled ovarian hyperstimulation (COH) is an effective method of assisted reproductive technology (ART) to stimulate the generation of more oocytes than are produced during natural cycles (1). Despite the increase in the number of embryos generated using COH, pregnancy rates following ART remain low at 20-30% per fresh cycle (2). Several studies and meta-analyses have shown that the gonadotropin-releasing

hormone analogues (GnRH-as), including GnRH agonists and antagonists, which are used in COH, may have negative effects on endometrial receptivity (3-5). However, the mechanisms regulating endometrial receptivity deficiency following GnRH-as treatment remain to be elucidated (2,4).

Homeobox (HOX) A10/Hoxa10 (human/mouse), respectively) is a homeobox-containing transcription factor that regulates embryo uterine development and is essential for endometrial development during each menstrual cycle in adults (6-7). Hoxa10 targeted mutation Hoxa10 (-/-) mice ovulate normally, but ~80% are sterile due to the low expression of maternal Hoxa10 in the distal oviductal and uterine epithelium, which may affect embryo implantation (8). Hoxa10 is a characteristic molecular marker of endometrial receptivity with peak expression exhibited during the window of implantation (9). Hoxa10 has been demonstrated to be involved in the regulation of pinopode development and downstream target genes, which are involved in implantation, including integrin β 3 (10-12).

Impaired endometrial receptivity is associated with altered Hoxa10 methylation. Abnormal expression of Hoxa10 has been found to be associated with disrupted endometrial receptivity in several diseases, including endometrial carcinoma, endometriosis, endometrial polyps, ovarian cancer, polycystic ovary syndrome and in conditions associated with exposure to diethylstilbestrol and bisphenol-A (13-18). However, few studies have demonstrated the association between the uterine methylation status and alterations in endometrial receptivity following GnRH-as therapies. The present study aimed to investigate the uterine methylation status of the Hoxa10 gene following GnRH-as treatment in order to explore the potential mechanism underlying the epigenetic effect of GnRH-as on endometrial receptivity.

Materials and methods

Animals. Animal care and use was conducted according to the institutional guidelines established by the Animal Care and Use Committee of Wuhan University (Wuhan, China) of 2010 Zhongnan Hospital of Wuhan University Animal Care and Use 025. Female, virgin BALB/c mice (7-9 weeks) were purchased from Hubei Medical Laboratory Animal Center (Wuhan, China) and were housed under a 12/12 h light/dark cycle at $25\pm 0.5^{\circ}\text{C}$ and 50-60% humidity. Mice were fed with

Correspondence to: Professor Yuanzhen Zhang, Department of Obstetrics and Gynecology, Reproductive Medicine Center, Zhongnan Hospital of Wuhan University, 169 Donghu Road, Wuhan, Hubei 430071, P.R. China
E-mail: 1151877287@qq.com

Key words: DNA methylation, endometrial receptivity, Hoxa10, gonadotropin-releasing hormone agonist, gonadotropin-releasing hormone analogue, gonadotropin-releasing hormone antagonist

a standard pellet diet and water. Smear samples of vaginal discharge were observed daily in order to identify the estrus. Only mice with more than two consecutive periods of regular 4-day estrus cycles were used in the present study. Suitable mice [age, 8-12 weeks; body weight (bw), 20-23 g] were randomly divided into three groups: The GnRH agonist treatment group (n=10), the GnRH antagonist treatment group (n=10) and the control (natural cycle) group (n=10).

Ovarian stimulation. Treatment procedures were performed as previously described, but with minor revisions (19). In brief, mice in the GnRH agonist group received intraperitoneal (IP) injection with the GnRH agonist Decapeptyl® (Ferring Co., Kiel, Germany) at 1.5 µg/100 g bw/day between days three and 9 of estrus. At 9 am of day 9, 20 IU/mouse human menopausal gonadotropin (HMG; Livzon Pharmaceutical Group Inc., Shanghai, China) was injected IP, followed by IP injection with 100 IU/100 g bw human chorionic gonadotrophin (HCG; Pregnyl®; Organon International, Oss, Netherlands) at 28 h after the injection of HMG. Mice in the GnRH antagonist group received IP injection of the GnRH antagonist Cetrotide® (Serono Inc., Rockland, MA, USA) at 4 µg/100 g bw on day three of estrus. HMG was then injected at 20 IU/mouse IP at 9 am of day 9, followed by IP injection with 100 IU/100 g bw HCG 28 h after the injection of HMG. The mice in the control (natural cycle) group received IP injection with saline only at the same volume as the injections received by the mice in the GnRH agonist and antagonist groups, from day three of estrus onwards. The same injection schedule was followed as described for the GnRH agonist and antagonist groups.

Tissue collection and application. Fresh whole uterus samples were collected from the mice in the three groups 48 h after the treatment. Fresh whole uterus samples were quickly divided into four equal sections subsequent to being washed in cold phosphate-buffered saline. One uterus section was fixed in 2.5% glutaraldehyde at room temperature for 30 min and at incubated at 4°C overnight, prior to being fixed for ≥1 h in 1% osmium tetroxide in the dark for scanning electron microscopy (SEM) analysis. The remaining three sections of each sample were stored at -80°C until required for protein, DNA and mRNA extraction.

Genomic DNA extraction and bisulfite sequencing polymerase chain reaction (BSP). Genomic DNA was extracted from the frozen tissue samples from the three groups using the DNeasy® Blood and Tissue kit (Qiagen, Valencia, CA, USA). The genomic DNA (500 ng) was then bisulfite-modified using the EZ DNA Methylation-Gold™ kit (Qiagen) according to the manufacturer's instructions. Bisulfite-modified DNA was dissolved in 20 µl water and stored at -80°C.

Quantification of Hoxa10 promoter methylation in mice using BSP. A total of 200 ng bisulfite-treated DNA was used in a 50 µl reaction system containing 1.5 µl forward and reverse primers (Table I), 1.25 mmol/l deoxynucleotide triphosphates, 25 mM Mg²⁺ and 0.5 µl HotStarTaq DNA polymerase (Qiagen). The amplification conditions were as follows: 10 min at 95°C followed by 40 cycles of 95°C for 30 sec, 53°C for 30 sec and 72°C for 40 sec, then a final extension at 72°C

for 10 min. Polymerase chain reaction (PCR) products were resolved using electrophoresis on a 2% agarose gel and stained with GoldView (SBS Genetech Co., Ltd., Beijing, China). The appropriate-sized product bands were then isolated and excised from the gel and purified using a Gel Extraction kit (Axygen Biotechnology, Hangzhou, China) according to the manufacturer's instructions. The resultant products were sequenced using MicoRead Biotechnology (Beijing Microread Genetics Co., Ltd., Beijing, China).

Quantitative polymerase chain reaction (qPCR) analysis. Total RNA was extracted from the tissue samples using the REzol RNA extraction kit (SBS Genetech Co., Ltd.) according to the manufacturer's instructions. Total RNA (100 ng) from each sample was treated with DNase and converted to complementary (c)DNA using the PrimeScript™ RT reagent kit (Takara Bio Inc., Dalian, China). mRNA levels were analyzed using an iQ5 Real-Time PCR detection system (Bio-Rad Laboratories, Inc., Hercules, CA, USA) with SYBR® Premix Ex Taq™ II (Takara Bio Inc.). The primer sequences for Hoxa10, integrin β3 and β-actin are listed in Table I. All primers were obtained from Servicebio (Wuhan, China).

The qPCR amplification conditions for Hoxa10 were as follows: 40 cycles of 95°C for 10 sec, 62°C for 15 sec and 72°C for 15 sec. The qPCR amplification conditions for integrin β3 and β-actin were as follows: 40 cycles of 95°C for 10 sec, 60°C for 15 sec and 72°C for 15 sec. The increasing fluorescence of the PCR products during amplification was monitored to create a quantitative standard curve. Quantification of the target gene expression in the samples was assessed and adjusted to the quantitative expression of β-actin in the same samples. Melting curve analysis was conducted to determine the specificity of the amplified products and to ensure the absence of primer-dimer formation. All products obtained yielded the predicted melting temperature. Relative gene expression was calculated using the 2^{-ΔΔCt} method.

Western blot analysis. Samples were lysed in radio-immunoprecipitation assay buffer (Beyotime, Shanghai, China) with protease inhibitors (Sigma-Aldrich, St. Louis, MO, USA) to extract whole-cell proteins. Protein levels were measured using the Micro Bicinchoninic Acid™ Protein Assay kit (Beyotime). Proteins were run on a precast 7.5% acrylamide gel (Beyotime) and transferred onto polyvinylidene fluoride membranes (Millipore Corporation, Billerica, MA, USA). Membranes were blocked with 5% fat-free milk in Tris-buffered saline with 0.1% Tween-20 followed by incubation with primary antibodies against Hoxa10 (AV100932; Sigma-Aldrich) and integrin β3 (ab33171; Abcam PLC, Cambridge, UK). Membranes were then incubated with secondary antibodies rabbit anti-goat polyclonal antibody (ZDR-5308; Beijing Zhongshan Golden Bridge Biotechnology Co., Ltd., Beijing, China) or goat anti-rabbit polyclonal antibody (ZDR-5306; Beijing Zhongshan Golden Bridge Biotechnology Co., Ltd.) in blocking buffer, respectively. Immunoreactive bands were detected using a chemiluminescent detection kit (Beyotime). Densitometry measurements were analyzed using Quantity One v4.4.0 (Bio-Rad Laboratories, Inc.). Target protein expression levels were normalized to that of β-actin.

Table I. Primer sequences used for BSP and qPCR.

Gene	Sense primer (5'-3')	Antisense primer (5'-3')
Hoxa10 (BSP)	TATTTTGAGGTAGTTTTTATAGTTT	CAAATAACCCCTTTCTAACTAACATTTC
Hoxa10 (qPCR)	CCTTCCGAGAGCAGCAAA	GTCTGGTGCTTCGTGTAGGG
Integrin β 3 (qPCR)	GCCTTCGTGGACAAGCCTGTA	GGACAATGCCTGCCAGTCTTC
β -actin (qPCR)	TTCCAGCCTTCCTTCCTGG	TTGCGCTCAGGAGGAGCAAT

Hox, homeobox; qPCR, quantitative polymerase chain reaction; BSP, Bisulfite sequencing PCR.




Figure 1. Methylation levels of the Hoxa10 gene promoter region in the uterus during the implantation window of GnRH analogue-exposed mice. ^aP<0.01 vs. the control (natural cycle) group; ^bP<0.01 vs. the GnRH antagonist group. Hox, homeobox; GnRH, gonadotropin-releasing hormone.

SEM analysis. SEM was performed for morphological analysis and to confirm the presence of pinopodes in the endometrium of mice during the implantation window in the natural cycle or following GnRH agonist or antagonist treatment. For SEM preparation, endometrial tissues were fixed in 2.5% glutaraldehyde at room temperature for 30 min then at 4°C overnight, prior to being fixed for at least 1 h in 1% osmium tetroxide in the dark. Samples were then dehydrated in a graded series of ethanol, critical-point-dried, mounted and coated with gold in a sputter coater (JFC-1300 Auto Fine Coater; Jeol Ltd., Tokyo, Japan). Samples were observed using a scanning electron microscope (JSM-5600LV SEM, Jeol Ltd.). SEM was performed in order to observe the morphology of the pinopodes in the samples from the different groups.

Statistical analysis. Statistical analyses for western blot analysis, BSP and qPCR values were performed using SPSS 17.0 software (SPSS, Inc., Chicago, IL, USA). χ^2 and analysis of variance tests were performed to compare the groups. P<0.05 was considered to indicate a statistically significant difference.

Results

Hoxa10 promoter methylation status and GnRH-as. BSP was performed to examine the methylation status of the 21 CpG

sites in the Hoxa10 promoter in each sample. In total, 210 CpG sites were analyzed in each experimental group. A total of 37 (17.6%) methylated CpG sites were observed in the samples from the GnRH antagonist group, compared with 14 (6.7%) methylated sites in the samples from the GnRH agonist group and four (1.9%) methylated sites in the samples from the control group (GnRH antagonist vs. control, P<0.001; GnRH agonist vs. control, P=0.006 and GnRH agonist vs. GnRH antagonist, P=0.008; Fig. 1). Thus, compared with the mice in the control group, the levels of methylation within the Hoxa10 promoter sequence were found to be higher in the mice that had received GnRH-as treatment, particularly in those treated with the GnRH antagonist.

Endometrial Hoxa10 and integrin β 3 mRNA expression during the implantation window following GnRH-as treatment. Hoxa10 and integrin β 3 mRNA were expressed during the implantation window in the uteri of the mice in all three groups. Following normalization to β -actin expression, Hoxa10 mRNA expression was observed to be significantly decreased in the mice in the GnRH antagonist and agonist groups compared with the control group (P<0.001 and P=0.004, respectively; Fig. 2A). Compared with the control group, integrin β 3 mRNA expression was also found to be reduced in the mice in the GnRH antagonist and agonist groups (P<0.001 and P=0.002, respectively; Fig. 2B). Furthermore, Hoxa10 and integrin β 3 mRNA expression were observed to be higher in the GnRH agonist group compared with the GnRH antagonist group (P=0.044 and P=0.032, respectively; Fig. 2A and B).

Endometrial Hoxa10 and integrin β 3 protein expression during the implantation window following GnRH-as treatment. The protein expression of endometrial Hoxa10 and integrin β 3 were detected using western blot analysis during the implantation window in the uteri of the mice in the three groups (Fig. 3A). In accordance with the mRNA expression findings, following normalization to β -actin expression, Hoxa10 protein expression was found to be lowest in the mice in the GnRH antagonist treatment group and highest in the mice in the control group (GnRH agonist vs. control, P=0.032; GnRH antagonist vs. control, P=0.047 and GnRH agonist vs. GnRH antagonist, P=0.005). Integrin β 3 expression was also observed to be lowest in the mice in the GnRH antagonist group and highest in those in the control group (GnRH agonist vs. control, P=0.006; GnRH antagonist vs. control P=0.004 and GnRH agonist vs. GnRH antagonist, P=0.0041; Fig. 3C).




Figure 2. Levels of (A) Hoxa10 and (B) integrin β 3 mRNA during the implantation window in the uteri of mice in the GnRH agonist, GnRH antagonist and control groups, measured using quantitative polymerase chain reaction analysis. Data are presented as the mean \pm standard error of the mean of the protein expression of the target gene relative to that of β -actin. The mRNA levels of Hoxa10 and integrin β 3 in the mice in the GnRH analogue groups, particularly the GnRH antagonist group, were lower than those in the control mice. ^{a,c}P<0.01 vs. control; ^{b,d}P<0.05 vs. GnRH agonist. Hox, homeobox; GnRH, gonadotropin-releasing hormone.




Figure 3. (A) Hoxa10 and integrin β 3 protein expression during the implantation window in the uteri of GnRH analogue-exposed mice, measured using western blot analysis. β -actin protein expression was used as an internal loading control. (B) Hoxa10 protein expression in the mice in the GnRH agonist, GnRH antagonist and control groups. (C) Integrin β 3 protein expression in the mice in the GnRH agonist, GnRH antagonist and control (natural cycle) groups. Data are presented as the mean \pm standard error of the mean of the protein expression of the target gene relative to that of β -actin. ^aP<0.05 vs. the control group; ^bP<0.01 vs. the GnRH agonist group; ^cP<0.01 vs. the control group; and ^dP<0.05 vs. the GnRH agonist group. Hox, homeobox; GnRH, gonadotropin-releasing hormone.




Figure 4. Scanning electron microscopy showing pinopodes of the luminal uterine epithelium during the implantation window in GnRH analogue-exposed mice. Mice in the GnRH agonist and GnRH antagonist groups exhibited reduced pinopode formation. (A) The luminal uterine epithelium of the mice in the control (natural cycle) group exhibited several fully developed pinopodes and a smooth surface; (B) the luminal uterine epithelium of the mice in the GnRH agonist group exhibited several developing pinopodes with a villous surface. (C) The luminal uterine epithelium of the mice in the GnRH antagonist group exhibited regressing pinopodes and a ciliated, microvillous surface with few/no pinopodes. Magnification, x5,000. GnRH, gonadotropin-releasing hormone.

Pinopode development following GnRH-as treatment. SEM photomicrographs of the endometrial luminal surface of the uterus during the implantation window were captured to

identify the developmental status of the pinopodes (Fig. 4). Pinopode development was observed to be repressed in the GnRH agonist and GnRH antagonist groups compared with

the control group. The mice in the control group exhibited fully developed pinopodes on the apical pole of the majority of the non-ciliated epithelial cells (Fig. 4A). However, in the mice in the GnRH agonist group, the pinopodes were less developed and the luminal uterine epithelium exhibited a villous surface (Fig. 4B). Pinopode development in the mice in the GnRH antagonist group was repressed and delayed, with the luminal uterine epithelium being ciliated and microvillous and exhibiting few pinopodes (Fig. 4C).

Discussion

The lower pregnancy rates in COH cycles during ART treatment may be a consequence of the negative effects of GnRH-as on endometrial receptivity; however, the specific mechanism has yet to be elucidated (1-2,20). The present study investigated Hoxa10 DNA methylation patterns and expression, as well as the effect of GnRH agonist and antagonist treatment on endometrial receptivity during the implantation window in mice.

In the present study, methylation in the promoter region of Hoxa10 was found to increase following GnRH agonist and antagonist treatment, with such increases being most evident under GnRH antagonist treatment. Hoxa10 is a well established biomarker for endometrial receptivity and alterations in its methylation have been demonstrated to be associated with disturbances in endometrial receptivity in several pathological endometrial conditions, including reproductive system diseases and exposure to environmental endocrine disruptors (12-17). Several studies have shown that methylation regulation may be involved in endometrial development during the adult menstrual cycle and endometrial decidualization (21-23). In addition, increasing evidence has suggested that epigenetic mechanisms may regulate numerous aspects of pregnancy and the outcome of ART, with roles in implantation, placentation and foetal growth (24-26). Therefore, Hoxa10 methylation may be involved in endometrial receptivity following GnRH-as treatment. To the best of our knowledge, this is the first study to show the methylation pattern of Hoxa10 following GnRH agonist and antagonist treatment.

In the present study, compared with the natural cycle control mice, Hoxa10 mRNA and protein expression was observed to decrease following GnRH-as treatment, particularly following GnRH antagonist treatment. This finding is supported by several studies, which have reported that despite there being no evidence to suggest that GnRH-as negatively affects oocyte quality, fertilization rates or embryo quality, GnRH-as treatment is associated with a statistically significant reduction in pregnancy rates (1-2). Thus, this reduction in pregnancy rates associated with GnRH-as treatment may be due to the negative effects associated with GnRH-as, particularly GnRH antagonists, on the endometrium and the repression of endometrial receptivity (27-29). Altered HOXA10 expression may be caused by aberrant methylation of the gene, with promoter hypermethylation often correlated with suppressed gene expression (9,30). Therefore, the aberrant expression of Hoxa10 in the mouse endometrium following GnRH-as treatment may result from altered Hoxa10 DNA methylation.

Like Hoxa10 expression, following GnRH-as treatment the expression of integrin β 3 and pinopode development were also reduced in the present study. Furthermore, Quinn and Casper (31) also reported that downregulation of Hoxa10 caused a reduced number of pinopodes, while overexpression of Hoxa10 resulted in an increase in pinopode numbers (31). Endometrial integrin β 3 and pinopodes are characteristic biomarkers closely associated with endometrial development and maturation and they peak in expression during the implantation window (31-33). Therefore, the repressed endometrial receptivity observed following GnRH-as treatment may be due to altered Hoxa10 expression. The findings of the present study are supported by previous studies reporting decreased integrin β 3 subunit expression following GnRH-as intervention (1,19,32).

Further investigations are required to identify the mechanism by which GnRH agonists and antagonists cause Hoxa10 gene promoter hypermethylation. The methylation pattern in humans may be different to that in mice; therefore, investigations on endometrial biopsies from patients undergoing COH should be performed in the future.

In conclusion, the present study has shown that GnRH-as may influence the methylation status of the Hoxa10 gene in mice, which may affect uterine receptivity and repress the expression of endometrial integrin β 3 and pinopode development. These findings present a potential epigenetic mechanism by which GnRH-as, particularly GnRH antagonists, may negatively affect endometrial receptivity. These findings may explain the low implantation rate associated with COH treatments, which involve GnRH-as in human *in vitro* fertilization clinics. However, additional studies are required to analyze the impact of methylation regulation, particularly epigenetic regulation by GnRH-as, on the endometrium.

Acknowledgements

The present study was supported by grants from the National Key Basic Research Program of China (grant no. 2012CB526600), the Key Research Program from the Health Department of Hubei Province (grant no. JX5A05), National Natural Science Foundation of China (grant no. 81100411) and the Fundamental Research Funds for the Central Universities of China (grant no. 2012303020204).

References

1. Rackow BW, Kliman HJ and Taylor HS: GnRH antagonists may affect endometrial receptivity. *Fertil Steril* 89: 1234-1239, 2008.
2. Al-Inany HG, Abou-Setta AM and Aboulghar M: Gonadotrophin-releasing hormone antagonists for assisted conception: a Cochrane review. *Reprod Biomed Online* 14: 640-649, 2007.
3. Bourgain C and Devroey P: The endometrium in stimulated cycles for IVF. *Hum Reprod Update* 9: 515-522, 2003.
4. Devroey P, Bourgain C, Macklon NS and Fauser BC: Reproductive biology and IVF: ovarian stimulation and endometrial receptivity. *Trends Endocrinol Metab* 15: 84-90, 2004.
5. Kolibianakis E, Bourgain C, Albano C, *et al*: Effect of ovarian stimulation with recombinant follicle-stimulating hormone, gonadotropin releasing hormone antagonists, and human chorionic gonadotropin on endometrial maturation on the day of oocyte pick-up. *Fertil Steril* 78: 1025-1029, 2002.
6. Taylor HS: The role of HOX genes in human implantation. *Hum Reprod Update* 6: 75-79, 2000.

7. Taylor HS, Vanden Heuvel GB and Igarashi P: A conserved HOX axis in the mouse and human female reproductive system: late establishment and persistent adult expression of the HOXA cluster genes. *Biol Reprod* 57: 1338-1345, 1997.
8. Satokata I, Benson G and Maas R: Sexually dimorphic sterility phenotypes in HOXA10-deficient mice. *Nature* 374: 460-463, 1995.
9. Wu Y, Halverson G, Basir Z, *et al*: Aberrant methylation at HOXA10 may be responsible for its aberrant expression in the endometrium of patients with endometriosis. *Am J Obstet Gynecol* 193: 371-380, 2005.
10. Daftary GS, Troy PJ, Bagot CN, *et al*: Direct regulation of beta3-integrin subunit gene expression by HOXA10 in endometrial cells. *Mol Endocrinol* 16: 571-579, 2002.
11. Taylor HS, Daftary GS and Selam B: Endometrial HOXA10 expression after controlled ovarian hyperstimulation with recombinant follicle-stimulating hormone. *Fertil Steril* 80 (Suppl 2): 839-843, 2003.
12. Kim JJ, Taylor HS, Lu Z, *et al*: Altered expression of HOXA10 in endometriosis: potential role in decidualization. *Mol Hum Reprod* 13: 323-332, 2007.
13. Nasu K, Kawano Y, Tsukamoto Y, *et al*: Aberrant DNA methylation status of endometriosis: epigenetics as the pathogenesis, biomarker and therapeutic target. *J Obstet Gynaecol Res* 37: 683-695, 2011.
14. Rackow BW, Jorgensen E and Taylor HS: Endometrial polyps affect uterine receptivity. *Fertil Steril* 95: 2690-2692, 2011.
15. Widschwendter M, Apostolidou S, Jones AA, *et al*: HOXA methylation in normal endometrium from premenopausal women is associated with the presence of ovarian cancer: a proof of principle study. *Int J Cancer* 125: 2214-2218, 2009.
16. Bromer JG, Wu J, Zhou Y and Taylor HS: Hypermethylation of homeobox A10 by in utero diethylstilbestrol exposure: an epigenetic mechanism for altered developmental programming. *Endocrinology* 150: 3376-3382, 2009.
17. Varayoud J, Ramos JG, Bosquiaz VL, Muñoz-de-Toro M, and Luque EH: Developmental exposure to Bisphenol a impairs the uterine response to ovarian steroids in the adult. *Endocrinology* 149: 5848-5860, 2008.
18. Hiyama M, Choi EK, Wakitani S, *et al*: Bisphenol-A (BPA) affects reproductive formation across generations in mice. *J Vet Med Sci* 73: 1211-1215, 2011.
19. Ruan HC, Zhu XM, Luo Q, *et al*: Ovarian stimulation with GnRH agonist, but not GnRH antagonist, partially restores the expression of endometrial integrin beta3 and leukaemia-inhibitory factor and improves uterine receptivity in mice. *Hum Reprod* 21: 2521-2529, 2006.
20. Shapiro BS, Daneshmand ST, Garner FC, *et al*: Evidence of impaired endometrial receptivity after ovarian stimulation for in vitro fertilization: a prospective randomized trial comparing fresh and frozen-thawed embryo transfer. *Fertil Steril* 96: 344-348, 2011.
21. Yamagata Y, Asada H, Tamura I, *et al*: DNA methyltransferase expression in the human endometrium: down-regulation by progesterone and estrogen. *Hum Reprod* 24: 1126-1132, 2009.
22. Munro SK, Farquhar CM, Mitchell MD and Ponnampalam AP: Epigenetic regulation of endometrium during the menstrual cycle. *Mol Hum Reprod* 16: 297-310, 2010.
23. Vincent ZL, Farquhar CM, Mitchell MD and Ponnampalam AP: Expression and regulation of DNA methyltransferases in human endometrium. *Fertil Steril* 95: 1522-1525, 2011.
24. Logan PC, Ponnampalam AP, Rahnama F, *et al*: The effect of DNA methylation inhibitor 5-Aza-2'-deoxycytidine on human endometrial stromal cells. *Hum Reprod* 25: 2859-2869, 2010.
25. van Montfoort AP, Hanssen LL, de Sutter P, *et al*: Assisted reproduction treatment and epigenetic inheritance. *Hum Reprod Update* 18: 171-197, 2012.
26. Horsthemke B and Ludwig M: Assisted reproduction: the epigenetic perspective. *Hum Reprod Update* 11: 473-482, 2005.
27. Tarlatzis BC and Bili HN: Gonadotropin-releasing hormone antagonists: im-pact of IVF practice and potential nonas-sisted reproductive technology applications. *Curr Opin Obstet Gynecol* 15: 259-264, 2003.
28. Ludwig M, Katalinic A and Diedrich K: Use of GnRH antagonists in ovarian stimulation for assisted reproductive technologies compared to the long protocol. *Meta-analysis. Arch Gynecol Obstet* 265: 175-182, 2001.
29. Olivennes F, Cunha-Filho JS, Fanchin R, *et al*: The use of GnRH antagonists in ovarian stimulation. *Hum Reprod Update* 8: 279-290, 2002.
30. Grimaldi G, Christian M, Quenby S and Brosens JJ: Expression of epigenetic effectors in decidualizing human endometrial stromal cells. *Mol Hum Reprod* 18: 451-458, 2012.
31. Quinn CE and Casper RF: Pinopodes: a questionable role in endometrial receptivity. *Hum Reprod Update* 15: 229-236, 2009.
32. Creus M, Ordi J, Fábregues F, *et al*: $\alpha\beta3$ integrin expression and pinopod formation in normal and out-of-phase endometria of fertile and infertile women. *Hum Reprod* 17: 2279-2286, 2002.
33. Creus M, Ordi J, Fábregues F, *et al*: The effect of different hormone therapies on integrin expression and pinopode formation in the human endometrium: a controlled study. *Hum Reprod* 18: 683-693, 2003.